LIST OF TABLES

Table		Page
1.	Scientific and common names for fish of known or probable occurrence in the Roanoke River and Coniot Creek in the vicinity of Company Swamp, Bertie County, North Carolina	35
2.	NPDES dischargers to the lower Roanoke River Basin	37
3.	Descriptions of the fixed sampling locations used in the striped bass food and feeding study, 1982-1988	38
4.	Taxonomic relationships of zooplankton collected from the lower Roanoke River, delta, and western Albemarle Sound, North Carolina, 1984-1991	42
5.	Mean annual and second quarter (April-June) instream flows (cfs) ± standard error (S.E.), of the lower Roanoke River, North Carolina, and number of days within specific flow regimes, 1984-1990 based on daily records of the USGS gage at Roanoke Rapids	45
6.	Normal and observed rainfall (inches) for the Roanoke River basin downstream of Kerr Reservoir (RM 178.7), and basinwide, for April-June 1963-1991	46
7.	Upstream and Roanoke River delta water quality comparisons for 1988 and 1990.	47
8.	Most frequently occurring phytoplankton taxa, and their relative occurrence in samples (%), in the lower Roanoke River and western Albemarle Sound, North Carolina, from spring 1984 through spring 1988	48
9.	Relative contribution (% using density) of each taxonomic group to the spring zooplankton community of the lower Roanoke River (Stations 1-12), North Carolina, 1984-1988	49
10.	Relative contribution (% using density) of each taxonomic group to the spring zooplankton community of Batchelor Bay (Stations 13-16), North Carolina, 1984-1988	51
11.	Relative contribution (% using density) of each taxonomic group to the spring zooplankton community of Western Albemarle Sound (Stations 17-32), North Carolina, 1984-1988	53